


## PRACTICE TEST 2 FRQ #1

- Let  $R$  be the region bounded by the  $x$ -axis, the graph of  $y = \sqrt{x}$ , and the line  $x = 4$ .
  - Find the area of the region  $R$ .
  - Find the value of  $h$  such that the vertical line  $x = h$  divides the region  $R$  into two regions of equal area.
  - Find the volume of the solid generated when  $R$  is revolved about the  $x$ -axis.
  - The vertical line  $x = k$  divides the region  $R$  into two regions such that when these two regions are revolved about the  $x$ -axis, they generate solids with equal volumes. Find the value of  $k$ .



$$A = \int_0^4 \sqrt{x} \, dx = \frac{2}{3} x^{3/2} \Big|_0^4 = \frac{16}{3} \text{ or } 5.333$$

(b)

$$\int_0^h \sqrt{x} \, dx = \frac{8}{3} \quad \text{---or---} \quad \int_0^h \sqrt{x} \, dx = \int_h^4 \sqrt{x} \, dx$$

$$\frac{2}{3} h^{3/2} = \frac{8}{3} \quad \text{---or---} \quad \frac{2}{3} h^{3/2} = \frac{16}{3} - \frac{2}{3} h^{3/2}$$

$$h = \sqrt[3]{16} \text{ or } 2.520 \text{ or } 2.519$$

(c)

$$V = \pi \int_0^4 (\sqrt{x})^2 \, dx = \pi \frac{x^2}{2} \Big|_0^4 = 8\pi$$

or 25.133 or 25.132

(d)

$$\pi \int_0^k (\sqrt{x})^2 \, dx = 4\pi \quad \text{---or---} \quad \pi \int_0^k (\sqrt{x})^2 \, dx = \pi \int_k^4 (\sqrt{x})^2 \, dx$$

$$\pi \frac{k^2}{2} = 4\pi \quad \text{---or---} \quad \pi \frac{k^2}{2} = 8\pi - \pi \frac{k^2}{2}$$

$$k = \sqrt{8} \text{ or } 2.828$$

$$2 \left\{ \begin{array}{l} 1: A = \int_0^4 \sqrt{x} \, dx \\ 1: \text{answer} \end{array} \right.$$

$$2 \left\{ \begin{array}{l} 1: \text{equation in } h \\ 1: \text{answer} \end{array} \right.$$

$$3 \left\{ \begin{array}{l} 1: \text{limits and constant} \\ 1: \text{integrand} \\ 1: \text{answer} \end{array} \right.$$

$$2 \left\{ \begin{array}{l} 1: \text{equation in } k \\ 1: \text{answer} \end{array} \right.$$

## PRACTICE TEST 2 FRQ #2

2. Let  $f$  be the function given by  $f(x) = 2xe^{2x}$ .

(a) Find  $\lim_{x \rightarrow -\infty} f(x)$  and  $\lim_{x \rightarrow \infty} f(x)$ .

(b) Find the absolute minimum value of  $f$ . Justify that your answer is an absolute minimum.

(c) What is the range of  $f$ ?

(d) Consider the family of functions defined by  $y = bxe^{bx}$ , where  $b$  is a nonzero constant. Show that the absolute minimum value of  $bxe^{bx}$  is the same for all nonzero values of  $b$ .

(a)  $\lim_{x \rightarrow -\infty} 2xe^{2x} = 0$

$\lim_{x \rightarrow \infty} 2xe^{2x} = \infty$  or DNE

(b)  $f'(x) = 2e^{2x} + 2x \cdot 2 \cdot e^{2x} = 2e^{2x}(1 + 2x) = 0$

if  $x = -1/2$

$f(-1/2) = -1/e$  or  $-0.368$  or  $-0.367$

$-1/e$  is an absolute minimum value because:

(i)  $f'(x) < 0$  for all  $x < -1/2$  and  
 $f'(x) > 0$  for all  $x > -1/2$

–or–

(ii)  $f'(x) \quad \begin{array}{c} - \qquad \qquad + \\ \hline \qquad \qquad | \qquad \qquad \\ \qquad \qquad -1/2 \end{array}$

and  $x = -1/2$  is the only critical number

(c) Range of  $f = [-1/e, \infty)$

or  $[-0.367, \infty)$

or  $[-0.368, \infty)$

(d)  $y' = be^{bx} + b^2xe^{bx} = be^{bx}(1 + bx) = 0$

if  $x = -1/b$

At  $x = -1/b$ ,  $y = -1/e$

$y$  has an absolute minimum value of  $-1/e$  for all nonzero  $b$

**2**  $\left\{ \begin{array}{l} 1: 0 \text{ as } x \rightarrow -\infty \\ 1: \infty \text{ or DNE as } x \rightarrow \infty \end{array} \right.$

**3**  $\left\{ \begin{array}{l} 1: \text{ solves } f'(x) = 0 \\ 1: \text{ evaluates } f \text{ at student's critical point} \\ \quad 0/1 \text{ if not local minimum from} \\ \quad \text{student's derivative} \\ 1: \text{ justifies absolute minimum value} \\ \quad 0/1 \text{ for a local argument} \\ \quad 0/1 \text{ without explicit symbolic} \\ \quad \text{derivative} \end{array} \right.$

Note: 0/3 if no absolute minimum based on student's derivative


**1:** answer

Note: must include the left-hand endpoint; exclude the right-hand "endpoint"

**3**  $\left\{ \begin{array}{l} 1: \text{ sets } y' = be^{bx}(1 + bx) = 0 \\ 1: \text{ solves student's } y' = 0 \\ 1: \text{ evaluates } y \text{ at a critical number} \\ \quad \text{and gets a value independent of } b \end{array} \right.$

Note: 0/3 if only considering specific values of  $b$

### PRACTICE TEST 2 FRQ #3



$t$ (seconds)	$v(t)$ (feet per second)
0	0
5	12
10	20
15	30
20	55
25	70
30	78
35	81
40	75
45	60
50	72

3. The graph of the velocity  $v(t)$ , in ft/sec, of a car traveling on a straight road, for  $0 \leq t \leq 50$ , is shown above. A table of values for  $v(t)$ , at 5 second intervals of time  $t$ , is shown to the right of the graph.

- During what intervals of time is the acceleration of the car positive? Give a reason for your answer.
- Find the average acceleration of the car, in  $\text{ft}/\text{sec}^2$ , over the interval  $0 \leq t \leq 50$ .
- Find one approximation for the acceleration of the car, in  $\text{ft}/\text{sec}^2$ , at  $t = 40$ . Show the computations you used to arrive at your answer.
- Approximate  $\int_0^{50} v(t) dt$  with a Riemann sum, using the midpoints of five subintervals of equal length. Using correct units, explain the meaning of this integral.

(a) Acceleration is positive on  $(0, 35)$  and  $(45, 50)$  because the velocity  $v(t)$  is increasing on  $[0, 35]$  and  $[45, 50]$

3 { 1:  $(0, 35)$   
1:  $(45, 50)$   
1: reason

Note: ignore inclusion of endpoints

(b) Avg. Acc. =  $\frac{v(50) - v(0)}{50 - 0} = \frac{72 - 0}{50} = \frac{72}{50}$   
or  $1.44 \text{ ft}/\text{sec}^2$

1: answer

(c) Difference quotient; e.g.

$$\frac{v(45) - v(40)}{5} = \frac{60 - 75}{5} = -3 \text{ ft}/\text{sec}^2 \text{ or}$$

$$\frac{v(40) - v(35)}{5} = \frac{75 - 81}{5} = -\frac{6}{5} \text{ ft}/\text{sec}^2 \text{ or}$$

$$\frac{v(45) - v(35)}{10} = \frac{60 - 81}{10} = -\frac{21}{10} \text{ ft}/\text{sec}^2$$

–or–

Slope of tangent line, e.g.

$$\text{through } (35, 90) \text{ and } (40, 75): \frac{90 - 75}{35 - 40} = -3 \text{ ft}/\text{sec}^2$$

2 { 1: method  
1: answer

Note: 0/2 if first point not earned

(d)  $\int_0^{50} v(t) dt$   
 $\approx 10[v(5) + v(15) + v(25) + v(35) + v(45)]$   
 $= 10(12 + 30 + 70 + 81 + 60)$   
 $= 2530 \text{ feet}$

3 { 1: midpoint Riemann sum  
1: answer  
1: meaning of integral

This integral is the total distance traveled in feet over the time 0 to 50 seconds.

## PRACTICE TEST 2 FRQ #4

4. Let  $f$  be a function with  $f(1) = 4$  such that for all points  $(x, y)$  on the graph of  $f$  the slope is given by  $\frac{3x^2 + 1}{2y}$ .
- (a) Find the slope of the graph of  $f$  at the point where  $x = 1$ .
- (b) Write an equation for the line tangent to the graph of  $f$  at  $x = 1$  and use it to approximate  $f(1.2)$ .
- (c) Find  $f(x)$  by solving the separable differential equation  $\frac{dy}{dx} = \frac{3x^2 + 1}{2y}$  with the initial condition  $f(1) = 4$ .
- (d) Use your solution from part (c) to find  $f(1.2)$ .

(a)  $\frac{dy}{dx} = \frac{3x^2 + 1}{2y}$

$$\left. \frac{dy}{dx} \right|_{\substack{x=1 \\ y=4}} = \frac{3+1}{2 \cdot 4} = \frac{4}{8} = \frac{1}{2}$$

(b)  $y - 4 = \frac{1}{2}(x - 1)$

$$f(1.2) - 4 \approx \frac{1}{2}(1.2 - 1)$$

$$f(1.2) \approx 0.1 + 4 = 4.1$$

(c)  $2y \, dy = (3x^2 + 1) \, dx$

$$\int 2y \, dy = \int (3x^2 + 1) \, dx$$

$$y^2 = x^3 + x + C$$

$$4^2 = 1 + 1 + C$$

$$14 = C$$

$$y^2 = x^3 + x + 14$$

$$y = \sqrt{x^3 + x + 14} \text{ is branch with point } (1, 4)$$

$$f(x) = \sqrt{x^3 + x + 14}$$

(d)  $f(1.2) = \sqrt{1.2^3 + 1.2 + 14} \approx 4.114$

1: answer

2 { 1: equation of tangent line  
1: uses equation to approximate  $f(1.2)$

5 { 1: separates variables  
1: antiderivative of  $dy$  term  
1: antiderivative of  $dx$  term  
1: uses  $y = 4$  when  $x = 1$  to pick one function out of a family of functions  
1: solves for  $y$   
0/1 if solving a linear equation in  $y$   
0/1 if no constant of integration

Note: max 0/5 if no separation of variables

Note: max 1/5 [1-0-0-0-0] if substitutes value(s) for  $x$ ,  $y$ , or  $dy/dx$  before antidifferentiation

1: answer, from student's solution to the given differential equation in (c)


## PRACTICE TEST 2 FRQ #5

5. The temperature outside a house during a 24-hour period is given by

$$F(t) = 80 - 10 \cos\left(\frac{\pi t}{12}\right), \quad 0 \leq t \leq 24,$$

where  $F(t)$  is measured in degrees Fahrenheit and  $t$  is measured in hours.

- Sketch the graph of  $F$  on the grid below.
- Find the average temperature, to the nearest degree Fahrenheit, between  $t = 6$  and  $t = 14$ .
- An air conditioner cooled the house whenever the outside temperature was at or above 78 degrees Fahrenheit. For what values of  $t$  was the air conditioner cooling the house?
- The cost of cooling the house accumulates at the rate of \$0.05 per hour for each degree the outside temperature exceeds 78 degrees Fahrenheit. What was the total cost, to the nearest cent, to cool the house for this 24-hour period?



(b)

$$\begin{aligned} \text{Avg.} &= \frac{1}{14-6} \int_6^{14} \left[ 80 - 10 \cos\left(\frac{\pi t}{12}\right) \right] dt \\ &= \frac{1}{8} (697.2957795) \\ &= 87.162 \text{ or } 87.161 \\ &\approx 87^\circ \text{ F} \end{aligned}$$

(c)

$$\begin{aligned} \left[ 80 - 10 \cos\left(\frac{\pi t}{12}\right) \right] - 78 &\geq 0 \\ 2 - 10 \cos\left(\frac{\pi t}{12}\right) &\geq 0 \\ \left. \begin{array}{l} 5.230 \\ \text{or} \\ 5.231 \end{array} \right\} &\leq t \leq \left\{ \begin{array}{l} 18.769 \\ \text{or} \\ 18.770 \end{array} \right. \end{aligned}$$

(d)

$$\begin{aligned} C &= 0.05 \int_{\substack{5.231 \\ \text{or} \\ 5.230}}^{\substack{18.769 \\ \text{or} \\ 18.770}} \left( \left[ 80 - 10 \cos\left(\frac{\pi t}{12}\right) \right] - 78 \right) dt \\ &= 0.05(101.92741) = 5.096 \approx \$5.10 \end{aligned}$$

1: bell-shaped graph  
 minimum 70 at  $t = 0, t = 24$  only  
 maximum 90 at  $t = 12$  only

2: integral  
 1: limits and  $1/(14-6)$   
 1: integrand  
**3** { 1: answer  
 0/1 if integral not of the form  
 $\frac{1}{b-a} \int_a^b F(t) dt$

**2** { 1: inequality or equation  
 1: solutions with interval

2: integral  
 1: limits and 0.05  
 1: integrand  
**3** { 1: answer  
 0/1 if integral not of the form  
 $k \int_a^b (F(t) - 78) dt$

## PRACTICE TEST 2 FRQ #6

6. Consider the curve defined by  $2y^3 + 6x^2y - 12x^2 + 6y = 1$ .

- (a) Show that  $\frac{dy}{dx} = \frac{4x - 2xy}{x^2 + y^2 + 1}$ .
- (b) Write an equation of each horizontal tangent line to the curve.
- (c) The line through the origin with slope  $-1$  is tangent to the curve at point  $P$ . Find the  $x$ - and  $y$ -coordinates of point  $P$ .

(a)  $6y^2 \frac{dy}{dx} + 6x^2 \frac{dy}{dx} + 12xy - 24x + 6 \frac{dy}{dx} = 0$

$$\frac{dy}{dx}(6y^2 + 6x^2 + 6) = 24x - 12xy$$

$$\frac{dy}{dx} = \frac{24x - 12xy}{6x^2 + 6y^2 + 6} = \frac{4x - 2xy}{x^2 + y^2 + 1}$$

(b)  $\frac{dy}{dx} = 0$

$$4x - 2xy = 2x(2 - y) = 0$$

$$x = 0 \text{ or } y = 2$$

When  $x = 0$ ,  $2y^3 + 6y = 1$ ;  $y = 0.165$

There is no point on the curve with  $y$  coordinate of 2.

$y = 0.165$  is the equation of the only horizontal tangent line.

(c)  $y = -x$  is equation of the line.

$$2(-x)^3 + 6x^2(-x) - 12x^2 + 6(-x) = 1$$

$$-8x^3 - 12x^2 - 6x - 1 = 0$$

$$x = -1/2, \quad y = 1/2$$

–or–

$$\frac{dy}{dx} = -1$$

$$4x - 2xy = -x^2 - y^2 - 1$$

$$4x + 2x^2 = -x^2 - x^2 - 1$$

$$4x^2 + 4x + 1 = 0$$

$$x = -1/2, \quad y = 1/2$$

$$2 \left\{ \begin{array}{l} 1: \text{ implicit differentiation} \\ 1: \text{ verifies expression for } \frac{dy}{dx} \end{array} \right.$$

$$4 \left\{ \begin{array}{l} 1: \text{ sets } \frac{dy}{dx} = 0 \\ 1: \text{ solves } \frac{dy}{dx} = 0 \\ 1: \text{ uses solutions for } x \text{ to find equation of horizontal tangent lines} \\ 1: \text{ verifies which solutions for } y \text{ yield equations of horizontal tangent line} \end{array} \right.$$

Note: max 1/4 [1-0-0-0] if  $dy/dx = 0$  is not of the form  $g(x, y)/h(x, y) = 0$  with solutions for both  $x$  and  $y$

$$3 \left\{ \begin{array}{l} 1: \quad y = -x \\ 1: \text{ substitutes } y = -x \text{ into equation of curve} \\ 1: \text{ solves for } x \text{ and } y \end{array} \right.$$

–or–

$$3 \left\{ \begin{array}{l} 1: \text{ sets } \frac{dy}{dx} = -1 \\ 1: \text{ substitutes } y = -x \text{ into } \frac{dy}{dx} \\ 1: \text{ solves for } x \text{ and } y \end{array} \right.$$

Note: max 2/3 [1-1-0] if importing incorrect derivative from part (a)