

PRACTICE TEST #6

CALCULUS AB
SECTION I, Part A
Time—55 minutes
Number of questions—28

A CALCULATOR MAY NOT BE USED ON THIS PART OF THE EXAMINATION.

Directions: Solve each of the following problems, using the available space for scratchwork. After examining the form of the choices, decide which is the best of the choices given and fill in the corresponding oval on the answer sheet. No credit will be given for anything written in the test book. Do not spend too much time on any one problem.

In this test:

- (1) Unless otherwise specified, the domain of a function f is assumed to be the set of all real numbers x for which $f(x)$ is a real number.
- (2) The inverse of a trigonometric function f may be indicated using the inverse function notation f^{-1} or with the prefix “arc” (e.g., $\sin^{-1} x = \arcsin x$).

1. If $y = (x^3 + 1)^2$, then $\frac{dy}{dx} =$

(A) $(3x^2)^2$ (B) $2(x^3 + 1)$ (C) $2(3x^2 + 1)$ (D) $3x^2(x^3 + 1)$ (E) $6x^2(x^3 + 1)$

2. $\int_0^1 e^{-4x} dx =$

(A) $\frac{-e^{-4}}{4}$ (B) $-4e^{-4}$ (C) $e^{-4} - 1$ (D) $\frac{1}{4} - \frac{e^{-4}}{4}$ (E) $4 - 4e^{-4}$

3. For $x \geq 0$, the horizontal line $y = 2$ is an asymptote for the graph of the function f . Which of the following statements must be true?

- (A) $f(0) = 2$
- (B) $f(x) \neq 2$ for all $x \geq 0$
- (C) $f(2)$ is undefined.
- (D) $\lim_{x \rightarrow 2} f(x) = \infty$
- (E) $\lim_{x \rightarrow \infty} f(x) = 2$

4. If $y = \frac{2x + 3}{3x + 2}$, then $\frac{dy}{dx} =$


(A) $\frac{12x + 13}{(3x + 2)^2}$ (B) $\frac{12x - 13}{(3x + 2)^2}$ (C) $\frac{5}{(3x + 2)^2}$ (D) $\frac{-5}{(3x + 2)^2}$ (E) $\frac{2}{3}$

5. $\int_0^{\frac{\pi}{4}} \sin x \, dx =$

- (A) $-\frac{\sqrt{2}}{2}$ (B) $\frac{\sqrt{2}}{2}$ (C) $-\frac{\sqrt{2}}{2} - 1$ (D) $-\frac{\sqrt{2}}{2} + 1$ (E) $\frac{\sqrt{2}}{2} - 1$

6. $\lim_{x \rightarrow \infty} \frac{x^3 - 2x^2 + 3x - 4}{4x^3 - 3x^2 + 2x - 1} =$

- (A) 4 (B) 1 (C) $\frac{1}{4}$ (D) 0 (E) -1



Graph of f'

7. The graph of f' , the derivative of the function f , is shown above. Which of the following statements is true about f ?

- (A) f is decreasing for $-1 \leq x \leq 1$.
 (B) f is increasing for $-2 \leq x \leq 0$.
 (C) f is increasing for $1 \leq x \leq 2$.
 (D) f has a local minimum at $x = 0$.
 (E) f is not differentiable at $x = -1$ and $x = 1$.






8. $\int x^2 \cos(x^3) \, dx =$

- (A) $-\frac{1}{3} \sin(x^3) + C$
 (B) $\frac{1}{3} \sin(x^3) + C$
 (C) $-\frac{x^3}{3} \sin(x^3) + C$
 (D) $\frac{x^3}{3} \sin(x^3) + C$
 (E) $\frac{x^3}{3} \sin\left(\frac{x^4}{4}\right) + C$

9. If $f(x) = \ln(x + 4 + e^{-3x})$, then $f'(0)$ is

- (A) $-\frac{2}{5}$ (B) $\frac{1}{5}$ (C) $\frac{1}{4}$ (D) $\frac{2}{5}$ (E) nonexistent

10. The function f has the property that $f(x)$, $f'(x)$, and $f''(x)$ are negative for all real values x . Which of the following could be the graph of f ?




11. Using the substitution $u = 2x + 1$, $\int_0^2 \sqrt{2x+1} dx$ is equivalent to

- (A) $\frac{1}{2} \int_{-1/2}^{1/2} \sqrt{u} du$ (B) $\frac{1}{2} \int_0^2 \sqrt{u} du$ (C) $\frac{1}{2} \int_1^5 \sqrt{u} du$ (D) $\int_0^2 \sqrt{u} du$ (E) $\int_1^5 \sqrt{u} du$

12. The rate of change of the volume, V , of water in a tank with respect to time, t , is directly proportional to the square root of the volume. Which of the following is a differential equation that describes this relationship?

- (A) $V(t) = k\sqrt{t}$
 (B) $V(t) = k\sqrt{V}$
 (C) $\frac{dV}{dt} = k\sqrt{t}$
 (D) $\frac{dV}{dt} = \frac{k}{\sqrt{V}}$
 (E) $\frac{dV}{dt} = k\sqrt{V}$



Graph of f

13. The graph of a function f is shown above. At which value of x is f continuous, but not differentiable?

- (A) a (B) b (C) c (D) d (E) e

14. If $y = x^2 \sin 2x$, then $\frac{dy}{dx} =$

- (A) $2x \cos 2x$
 (B) $4x \cos 2x$
 (C) $2x(\sin 2x + \cos 2x)$
 (D) $2x(\sin 2x - x \cos 2x)$
 (E) $2x(\sin 2x + x \cos 2x)$

15. Let f be the function with derivative given by $f'(x) = x^2 - \frac{2}{x}$. On which of the following intervals is f decreasing?

- (A) $(-\infty, -1]$ only
 (B) $(-\infty, 0)$
 (C) $[-1, 0)$ only
 (D) $(0, \sqrt[3]{2}]$
 (E) $[\sqrt[3]{2}, \infty)$

x	-4	-3	-2	-1	0	1	2	3	4
$g'(x)$	2	3	0	-3	-2	-1	0	3	2

16. If the line tangent to the graph of the function f at the point $(1, 7)$ passes through the point $(-2, -2)$, then $f'(1)$ is

- (A) -5 (B) 1 (C) 3 (D) 7 (E) undefined

17. Let f be the function given by $f(x) = 2xe^x$. The graph of f is concave down when

- (A) $x < -2$ (B) $x > -2$ (C) $x < -1$ (D) $x > -1$ (E) $x < 0$


x	-4	-3	-2	-1	0	1	2	3	4
$g'(x)$	2	3	0	-3	-2	-1	0	3	2

18. The derivative g' of a function g is continuous and has exactly two zeros. Selected values of g' are given in the table above. If the domain of g is the set of all real numbers, then g is decreasing on which of the following intervals?

- (A) $-2 \leq x \leq 2$ only
 (B) $-1 \leq x \leq 1$ only
 (C) $x \geq -2$
 (D) $x \geq 2$ only
 (E) $x \leq -2$ or $x \geq 2$

19. A curve has slope $2x + 3$ at each point (x, y) on the curve. Which of the following is an equation for this curve if it passes through the point $(1, 2)$?


- (A) $y = 5x - 3$
- (B) $y = x^2 + 1$
- (C) $y = x^2 + 3x$
- (D) $y = x^2 + 3x - 2$
- (E) $y = 2x^2 + 3x - 3$



$$f(x) = \begin{cases} x + 2 & \text{if } x \leq 3 \\ 4x - 7 & \text{if } x > 3 \end{cases}$$


20. Let f be the function given above. Which of the following statements are true about f ?

- I. $\lim_{x \rightarrow 3} f(x)$ exists.
 - II. f is continuous at $x = 3$.
 - III. f is differentiable at $x = 3$.
- (A) None
 - (B) I only
 - (C) II only
 - (D) I and II only
 - (E) I, II, and III



21. The second derivative of the function f is given by $f''(x) = x(x - a)(x - b)^2$. The graph of f'' is shown above. For what values of x does the graph of f have a point of inflection?

- (A) 0 and a only
- (B) 0 and m only
- (C) b and j only
- (D) 0, a , and b
- (E) b , j , and k



22. The graph of f' , the derivative of f , is the line shown in the figure above. If $f(0) = 5$, then $f(1) =$


- (A) 0
- (B) 3
- (C) 6
- (D) 8
- (E) 11

23. $\frac{d}{dx} \left(\int_0^{x^2} \sin(t^3) dt \right) =$

- (A) $-\cos(x^6)$
- (B) $\sin(x^3)$
- (C) $\sin(x^6)$
- (D) $2x \sin(x^3)$
- (E) $2x \sin(x^6)$

24. Let f be the function defined by $f(x) = 4x^3 - 5x + 3$. Which of the following is an equation of the line tangent to the graph of f at the point where $x = -1$?

- (A) $y = 7x - 3$
- (B) $y = 7x + 7$
- (C) $y = 7x + 11$
- (D) $y = -5x - 1$
- (E) $y = -5x - 5$



25. A particle moves along the x -axis so that at time $t \geq 0$ its position is given by $x(t) = 2t^3 - 21t^2 + 72t - 53$. At what time t is the particle at rest?

- (A) $t = 1$ only
- (B) $t = 3$ only
- (C) $t = \frac{7}{2}$ only
- (D) $t = 3$ and $t = \frac{7}{2}$
- (E) $t = 3$ and $t = 4$

26. What is the slope of the line tangent to the curve $3y^2 - 2x^2 = 6 - 2xy$ at the point $(3, 2)$?

- (A) 0 (B) $\frac{4}{9}$ (C) $\frac{7}{9}$ (D) $\frac{6}{7}$ (E) $\frac{5}{3}$
-

27. Let f be the function defined by $f(x) = x^3 + x$. If $g(x) = f^{-1}(x)$ and $g(2) = 1$, what is the value of $g'(2)$?

- (A) $\frac{1}{13}$ (B) $\frac{1}{4}$ (C) $\frac{7}{4}$ (D) 4 (E) 13
-

28. Let g be a twice-differentiable function with $g'(x) > 0$ and $g''(x) > 0$ for all real numbers x , such that $g(4) = 12$ and $g(5) = 18$. Of the following, which is a possible value for $g(6)$?

- (A) 15 (B) 18 (C) 21 (D) 24 (E) 27
-