

## Basic facts about head lice



Head lice are small parasitic insects that do not come from out of the air or from the ground. They are human parasites and have probably been around since the beginning of time.

Head lice have been found on people of all socio-economic levels.

Personal hygiene or cleanliness in the home or school has nothing to do with getting head lice.

Children, particularly those of elementary school age, are most likely to get head lice because of their close contact with each other.

The main symptom is itching of the head.

There is no way to know where a child got head lice. Homes or schools do not get head lice – people do. Current research indicates that most lice infestations are rarely contracted in the school setting.

For more information:  
[www.headlice.org](http://www.headlice.org)



## **ISD Health Center Head Lice Policy**

Parents have prime responsibility of assisting in the prevention and management of head lice cases through regular checks of their child's hair and starting immediate treatment when head lice are detected.

While head lice are a nuisance, they do not spread disease and are not a health issue. Should a case of head lice be brought to the school's attention, the school nurse will maintain confidentiality at school, verify presence of an active infestation, and bring it to the family's attention. Parents, students and school staff will be educated about head lice identification, treatment and prevention.

If at all possible, students should **NOT** be excluded from school for having head lice, as the management of head lice should not disrupt the educational process of the child. The need to exclude students from school will be determined on a case by case basis.

This policy is supported by the National Pediculosis Association, the American Academy of Pediatrics, and the National Association of School Nurses.

**Indiana School for the Deaf  
Health Center**

**317-550-4818 (voice)  
317-614-0130 Fax number  
317-493-0497 VP**

**Deb Robarge  
Health Center Director  
[drobarge@isd.k12.in.us](mailto:drobarge@isd.k12.in.us)**

## What do head lice look like?

There are three forms of lice: the egg (also called a nit), the nymph, and the adult.

**Nits** are very small, about the size of a knot in thread, and hard to see.

They are oval in shape, and usually yellow to white in color. They are laid by the adult female at the base of the hair shaft and are firmly



attached with a glue-like substance. While nits take about one week to hatch, the ones that are likely to hatch are usually located within one fourth inch of the scalp. It is important not to confuse nits with dandruff, hair casts or other hair debris.

**Nymphs** are baby lice. They look like adult head lice, but are smaller in size. Nymphs mature into adults in about seven days after hatching. To live, the nymph must feed on human blood from the scalp.

**Adult lice** are about the size of a sesame seed and are tan to grayish white in color. Females, which are usually larger than males, lay 3-5 eggs a day. The eggs hatch in 7-10 days and it takes another 7-10 days for the louse to mature and lay their own eggs. Adult lice can live up to 30 days on a person's head. To live, adult lice need to feed on human blood from the scalp, behind the ears and on the back of the neck. They do not have hind legs or wings and cannot jump or fly, but they can crawl rapidly. If the louse falls off a person, it dies within 2 days.

## How are head lice spread?

Live head lice are spread through direct head to head contact with an infested person, but also by sharing hats, combs, and other hair accessories. By the time an infestation is found, the child is likely to have had it for at least a month. A child cannot catch nits, as nits can be laid only by live lice. Head lice are not spread by animals. Teach children not to share hats and grooming items (combs, hairbrushes, hair accessories, etc.) and to avoid direct head-to-head contact.



## Nit removal:

*The BEST treatment for removing Nits is to manually remove them. Complete nit removal is the key to successful head lice treatment.*

- Work in a well lighted area or use a flashlight and magnifying glass.
- Use a grooming comb or hairbrush to remove tangles. A hair de-tangle spray or other hair conditioner may aid in this process.. Divide the hair into sections and fasten off the hair that is not being worked on.
- Go through each hair section from the scalp to the end of the hair. Nits are usually found close to the scalp.
- Sift through the same section of hair and look for attached nits and live lice. Use your fingers to pull off the nits.
- Move on to the next section until the entire scalp and all hair has been checked. Screen the infested person every day for 10 days and regularly thereafter.

## Environmental clean up:

To kill lice on bedding, clothes, etc., wash in hot water and dry. Dry clean clothing that is not washable. Store clothing, stuffed animals, comforters, etc. that cannot be washed or dry cleaned into a plastic bag and seal for 2 weeks. Vacuum materials that cannot be washed (furniture, carpets). **Pesticide sprays are NOT recommended, as they have not been proven to be effective and they are a health hazard.** To kill lice on brushes, combs or hair accessories, soak them in hot water.

Contact the HC at ISD for more information.